




Hydrospatial Analysis

Homepage: www.gathacognition.com/journal/gcj3
<http://dx.doi.org/10.21523/gcj3>



Original Research Paper

Fluoride Accumulation in Groundwater from Semi-arid Part of Deccan Volcanic Province, India: A Cause of Urolithiasis Outbreak



Sanjay Kale^{1*}, N. J. Pawar²

1. Department of Environmental Sciences, Savitribai Phule Pune University, Pune - 411 007 (India).

2. Department of Geology, Savitribai Phule Pune University, Pune - 411 007 (India).

Abstract

Groundwater fluoride and health problem was meticulously studied for dental and skeleton fluorosis except few studies on urolithiasis. Urolithiasis is multi-factorial disease and excess fluoride consumption is one of the causal factors. In view of this, increase of fluoride in groundwater is reported in semiarid Deccan Volcanic Province (DVP), India. To understand the fluoride and urolithiasis association, present study was carried out in Karha river basin of DVP region. Three stages of data generation were adopted for present study such as procuring of medical records of urolithiasis, previous groundwater chemistry data and geochemical investigation of 50 groundwater samples from representative villages. Further, these variables were used for correlation analysis, temporal and spatial distribution to find out their relationships. Result shows medical records of hospitals indicating the gradual increase in urolithiasis is reported during drought situations. In temporal variation, annual fluoride concentration of groundwater and hot days are positively correlated with annual urolith patients as well as spatial study supports the same. In conclusion, present study highlights the relationship of urolith formation with number of hot days, groundwater electrical conductivity and fluoride. However, detailed biomedical study may lead towards understanding of fluoride- urolithiasis relationship.

Article history

Received: 02 August 2017

Accepted: 15 October 2017

Keywords

Fluoride;
Groundwater;
Deccan Volcanic Province;
Urolithiasis;
Maharashtra.

Editor(s)

P. S. Hire

© 2017 GATHA COGNITION® All rights reserved.

1 INTRODUCTION

Fluoride concentration of groundwater in India varies noticeably. Drought conditions admit less water in aquifer system accelerates the fluoride ion dissolution from parent lithology. The presence of the accessory minerals, fluorite and apatite (Handa, 1975), cryolite and topaz (Madhavan and Subramanian, 2001) in the rock mineral assemblage are the major source for dissolution of fluoride in groundwater. In addition, minerals like biotite, muscovite and hornblende may contain considerable percent of fluoride (Madhavan and Subramanian, 2001). The weathering of host rock as the results of monsoonal water is the major phenomenon for existence of ionic fluoride in groundwater. Its concentration is restricted to its trace occurrence in basaltic aquifers. Lowering of water table during pre-

monsoon period increases fluoride concentration in available water due to its high solubility. The consumption of fluoride through groundwater ingresses the human metabolism and chronic consumption leads to various health deformities like dental, skeleton fluorosis and nervous system impairment. In addition to this health hazard, renal system related illnesses are observed by some workers.

Averagely, 1 to 5% of world population facing urolith formation during their lifetime and is a most painful disorder of renal system (Sierakowski *et al.* 1979). Sudden increase in urolithiasis episode in specific areas due to dominance in arid climate is exerting tremendous pressure on medical management of urolithiasis. About 80% of uroliths are of calcium types

* Author address for correspondence

Department of Environmental Sciences, Savitribai Phule Pune University, Pune- 411 007 (India).

Tel.: +91 9890177791

E-mails: sanjakale@gmail.com (S. Kale-Corresponding author); njpawar@unipune.ac.in (N. Pawar).

<http://dx.doi.org/10.21523/gcj3.17010102>

© 2017 GATHA COGNITION® All rights reserved.

(calcium oxalate monohydrate/di-hydrate, calcium phosphate and mixture of all in varied proportion), 10 to 15% of uric acid stone, 1 to 5% struvite stone and rare stone of cystine stone (Asper, 1984). The majority of reasons for urolith formation are dietary habits, dehydration, hereditary, life style, old age prostate disorder and powerful medicine used in diseases like cancers and AIDS (Costa-Bauza *et al.*, 2007). However, increase in urolith patients in population suggest the role of environmental factors such as quality of water consumed as well as changing climatic conditions in urolith formation.

The renal system is responsible for excreting most of the body's excess fluoride and is exposed to higher concentrations of fluoride than are other organs (Whitford, 1996). This suggests that it might be at higher risk of fluoride toxicity than most soft tissues. Muralidharan *et al.* (2002) studied the high fluoride concentration found in arid regions of Rajasthan state in India and the conclusion was drawn as causative minerals like calcite and dolomite have accelerated the leaching of fluoride to the groundwater. It is to be noted that from north India the relationship between fluoride concentration in drinking water and urolith prevalence in local population is reported by Singh *et al.* (2001). The review article by Ozsvath (2009) highlighted the relationship between fluoride consumption through groundwater and its effects on renal system. However, inadequate information is available on occurrence of urolithiasis (kidney stone disease) from basaltic areas where groundwater is amassed by fluoride concentration during drought situations and its health implications in the form of urolithiasis.

Increased urolithiasis episodes were observed in semi-arid Karha river basin, Maharashtra from Deccan Volcanic Province (DVP) of peninsular India during drought period and exact reason is quite obscure for medical professionals as the majority of urolithiasis in the area is neither having genetic origin nor anatomical reason. It necessitates understanding the possibility of environmental causative factors for urolith formation in the study area.

For this intend present study was undertaken to identify the urolithiasis occurrence with respect to consumption of highly mineralized groundwater.

2 STUDY AREA

The area under investigation is Karha river basin (Figure 1) with a geographical area of about 1333 km², from semiarid region of Deccan Volcanic Province (DVP). The Karha basin has geographical coordinates of 73° 52' 07" E to 74° 41' 01" E and 18° 3' 41" N to 18° 26' 31" N. The major settlements of study area are Saswad and Baramati.

Study area present in semiarid and rain fed region of Maharashtra. In year 2000 to 2004 drought situations were experienced due to declined annual rainfall < 550 mm. Such drought conditions developed the severe water scarcity in the region, which prompted people to depend upon highly saline water supply for drinking purpose.

Groundwater withdrawal is confined to vesicular, weathered, jointed and fractured upper basaltic crust, which is overlain by thin soil cover. Dug wells are the principal source of drinking water supply in the study area having average depth up to 10 m. Depleting groundwater levels are common in pre-monsoon season and condition being further aggravated by drought situations. Average groundwater level during pre-monsoon is 8 m whereas the water tables are fairly shallower during post-monsoon with an average depth of 4.6 m. The erratic nature of south-west monsoon is the controlling factor for groundwater chemistry variation.

3 MATERIALS AND METHODS

To understand the temporal occurrence of urolithiasis from study area, previous 15 year data of hospitalized urolith patients was used. From this data, epidemic trends like number of urolith patients and gender class distribution were estimated. The information about number of 7081 urolith episodes reported in hospitals in the territory of study area (1994-2008) was procured for epidemiological study. Climatic data and groundwater quality data were acquired from government agencies namely IMD (India Meteorological Department) and GSDA (Groundwater Surveys and Development Authority), Pune, Government of Maharashtra. Number of summer hot days are calculated to their Tmax 97.5 > = percentile for 40.6°C (Gadgil and Dhorade, 2005).

Further, spatial (village-wise) variation of groundwater fluoride data was used to correlate with village wise urolithiasis occurrence. For this purpose, 50 groundwater samples were collected from representative villages (Figure 1) during two consecutive post-monsoon (December, 2004) and pre-monsoon (May 2005) seasons. Physicochemical analyses of collected groundwater samples were carried out by using standard methods (A.P.H.A., 1995). In that, fluoride from groundwater was determined with SPADNS method by using Nano-colorimeter (500D) (Kale *et al.*, 2010).

The data was generated for average groundwater chemistry and urolith patient reported during respective years. Annual pattern of groundwater fluoride concentration and annual urolithiasis occurrence is correlated for temporal study. However, village wise occurrence of urolithiasis is used to correlate with respective groundwater fluoride concentration of dug well respective village.




Figure 1. Location map and water sampling station of Karha basin in Deccan Volcanic Province (DVP), India

4 RESULTS

The results are shown in following two subheadings: 1) Epidemiology of urolithiasis 2) Intra-annual climatic variation and urolithiasis 3) Intra-annual variations in groundwater chemistry and urolithiasis, 4) Spatial distribution of urolithiasis and groundwater chemistry

4.1 Epidemiology of Urolithiasis

The urolith patient data of 7081 was procured from local hospitals and processed for its gender class as well as its annual distribution. Increasing trends of urolith patients are observed from year 2000 and male population is

more prone to urolithiasis than female (Figure 2). The average male to female ratio of urolith is observed 2.88 (with SD 2.15) during period 1994 to 2008, indicating vulnerability of male patients for urolithiasis in the study area.

From medical records of hospitals more than 95% of uroliths are of calcium oxalate type and size ranges from 2 mm to 15 mm. However, few uroliths are bigger in sizes up to 75 mm in diameter are formed due to various anatomical disorders such as genetically originated, hyper calciuria generated, old age prostate




Figure 2. Intra-annual gender distribution of urolith patients from Karha basin (Kale *et al.* 2014)

disorder and miscellaneous due to consumption of higher drugs in various treatments. Tiny uroliths of calcium oxalate are formed in majority of urolith patients from study area and altered geo-environmental and climatic conditions in drought situation are the focus points of present study.

4.2 Intra-annual Climatic Variation and Urolithiasis

Climatic data and urolith patient data of year 1994 to 2008 were used for temporal study (Table 1). More than

500 mm rainfall is observed from year 1995 to 1999 and substantial decrease (< 500 mm) is observed from year 2000 to 2003 (4 year). However, rainfall is increased from year 2004 to 2007. Total annual rainfall is poorly correlated to number of urolith patients reported in respective year. In less rainfall year of 2005, 2006 and 2007, number of hot days are showing more than 5 days and there is substantial increase in number of hot days 12 and 10 during year 2007 and 2008 respectively (Table 1). Number of hot days in a year is positively

Table 1. Annual weather station data corresponding to Karha basin

Year	Total annual rainfall in mm	Annual number of hot days	Total annual pan evaporation in mm	Total annual sunshine hours
1994	398.55	0	1960.7	2672.3
1995	731.05	1	1947.5	2758.1
1996	937.5	0	1981.9	2772.7
1997	376.5	0	1909.9	2907.3
1998	1046	3	1865.9	2684.9
1999	621.5	2	1720.5	2612.8
2000	342.7	2	1862.0	2841.8
2001	482.75	5	1756.4	2717.8
2002	238.9	8	1880.7	2791.0
2003	164.1	5	1843.5	2792.8
2004	567.5	1	1855.1	2811.2
2005	930	3	1754.9	2649.5
2006	971.5	4	1772.1	2616.6
2007	596.5	12	1858.4	2629.5
2008	470.5	10	1765.1	2579.7

^a Rainfall is average values of Saswad and Baramati rain gauge stations, ^bTemperature, pan evaporation and sunshine hours are of IMD Pune as a nearest weather station to Karha basin. ^c Hot days are calculated at Tmax 37.5 > = percentile for 40.6° C (Kale *et al.* 2014).

correlated ($R^2 = 0.8322$) with urolith patients from study area. Generally, 1-3 hot days are experienced in study area per year but in drought conditions the number exceeds up to more than four.

4.3 Intra-annual Variations in Groundwater Chemistry and Urolithiasis

Annual average groundwater chemical analysis data of 24 observatory wells of GSDA in study area was used to correlate with annually reported urolith patient (Table 2). Positive correlation ($R^2 = 0.7588$) of average annual groundwater conductivity with annual reported urolith patients is observed from study area. Other groundwater ionic species such as Ca, Mg, Na, K, HCO_3 , Cl, SO_4 and NO_3 are showing meagre correlation with urolith episodes in the study area. Water types observed in year having rainfall > 500 mm is Ca+Mg- HCO_3 and water type of Na+K-Cl+ SO_4 is dominated in year having rainfall < 500 mm. However, in the case of F ion there is significantly positive correlation with urolithiasis ($R^2 = 0.85$).

4.4 Spatial Distribution of Urolithiasis and Groundwater Chemistry

Spatial (village-wise) distributions of urolithiasis cases (during years: 2004 and 2005) and fluoride concentration in respective groundwater source was studied to understand their relationship. In post-monsoon (year 2004), urolithiasis occurrence was observed from 29 villages, however in pre-monsoon 18 villages are affected by urolithiasis. This urolithiasis occurrence were calculated for its density per 10,000

individuals (Indridason *et al.*, 2006), however fluoride concentration is expressed in epm. In present study, maximum fluoride concentration observed in post-monsoon (December 2004) and pre-monsoon (May 2005) are 3.96 mg/L and 3.40 mg/L respectively. In post-monsoon season, village-wise density of urolith patients is convincingly correlated to groundwater fluoride concentration (Figure 3a). However, numbers of villages are decreased for urolith patient occurrence in pre-monsoon season and the positive correlation is observed with groundwater fluoride concentration (Figure 3b).

5 DISCUSSION

5.1 Epidemiology of Urolithiasis

During year 1994 to 1999 urolith patients are observed less than 100 patients per year. This is due the satisfactorily rainfall experiences more than 500 mm, however consecutive decrease in rainfall (< 500 mm) for next four years showing continuous increase in urolith patients up to year 2008. As the groundwater is only source available for drinking purposes in the study area and people have to fulfil their drinking water need with depleted water source.

However, consecutive drought year increases mineralization potential of groundwater from study area due to decline in water level and evaporation dominance in semi-arid climate. This indicates that though rainfall is higher in post drought years the urolithiasis occurrence is increasing substantially. The groundwater mineralization potential is influenced by rainfall of previous year and groundwater exploitation.

Table 2. Annual averages of groundwater quality parameters from Karha basin during period 1994 – 2008

Year	W. L.	E.C.	Ca	Mg	Na	K	HCO_3	Cl	SO_4	NO_3	F
1994	7.6	592.5	16.4	42.8	17.9	0.2	72.8	77.5	16.7	22.8	0.4
1995	7.5	676.9	54.1	38.6	22.1	0.2	124.6	170.0	47.0	13.4	1.1
1996	7.9	706.9	47.8	40.8	23.0	0.3	135.2	180.0	93.5	20.8	1.1
1997	7.5	745.0	25.3	25.9	22.8	0.4	130.0	115.0	37.7	23.0	0.5
1998	8.3	690.0	19.1	19.3	23.9	0.3	109.0	81.8	33.0	24.5	0.4
1999	8.1	687.5	22.8	22.9	28.9	0.7	101.5	84.3	31.0	26.0	0.5
2000	8.9	952.3	66.2	36.7	51.8	0.5	140.7	159.8	50.4	1.5	0.7
2001	11.1	1086.8	60.0	26.9	122.3	1.0	206.7	203.4	125.4	2.3	0.8
2002	12.0	1375.3	100.2	97.7	398.3	1.6	148.6	531.1	450.3	11.1	1.1
2003	17.0	1203.8	117.4	75.7	200.0	5.6	130.3	400.1	254.9	10.1	0.9
2004	18.0	999.7	104.0	50.2	173.7	6.4	99.5	272.5	94.0	3.2	0.8
2005	11.8	1112.2	55.8	29.9	103.2	6.6	162.0	158.9	96.6	16.8	1.3
2006	8.4	1797.3	44.1	58.8	202.8	7.8	328.0	226.0	131.8	18.8	1.5
2007	9.6	2015.9	76.3	60.2	323.0	5.2	202.0	284.7	349.4	26.1	2.9
2008	10.1	1283.0	56.6	42.8	231.8	3.9	170.0	217.5	219.0	20.6	2.3

^a All parameters are in mg/L except W. L. is groundwater level in meters and electrical conductivity (E. C.) is measured in $\mu\text{S}/\text{cm}$. (Kale *et al.* 2014)




Figure 3. Relation between of urolith patient density per 10,000 inhabitants with fluoride concentration in groundwater from Karha basin: a) post-monsoon 2004, b) pre-monsoon 2005

5.2 Intra-annual Climatic Variation and Urolithiasis

In results of climatic variation and urolithiasis is shown in section 4.2, in that annual rainfall is poorly correlated with annual incidences of urolithiasis. Consecutive drought years depleted the groundwater up to its critical level and even if good rainfall years after drought also shows more groundwater mineralization potential. This situation is deprives the correlation between rainfall and urolithiasis. Consecutive drought years shows the cumulative built up of ionic concentration in groundwater from study are as the result of less water available for flushing of accumulated salts in vadose zone.

Further, number of hot days is convincingly correlated with urolithiasis in the study area. The reason for this is the heat stroke occurs that time the body is unable to regulate the temperature which causes rise in body temperature (Loughnan *et al.*, 2010). Urinary tract stones in the regular season in summer heat caused are renal colic because of hot weather leading to substantial increase in human perspiration, resulting in severe loss of body water and reduce the urine volume (Chandrajith *et al.*, 2006). Therefore, in the case of stone patients or potential patients with induced renal colic stone is the result of concentrated urine and allied compounds (Hassan *et al.*, 2001) and finally results into urolith formation.

5.3 Basaltic Groundwater Quality Variation form Year 1994 to 2008

The water type of Ca+Mg-HCO₃ dominance is observed during year during year which experiences rainfall > 500 mm. However, water type is shifted towards Na+K-Cl+SO₄ during year experiencing rainfall < 500 mm as the results groundwater salinization due to depleting water table and evaporation dominance in vadose zone. The geochemical shifting of pristine alkaline aquifer condition to highly saline water dominated by Na+K-Cl+SO₄ constituents.

In basaltic groundwater, ionic dissolution is predominantly derived from weathering phenomenon of ferromagnetic silicates (olivine and augite) and plagioclase feldspar. Also some ionic contribution is possible by dissolution of secondary minerals like zeolites. It is seen that fluoride concentration in groundwater correlates positively with urolith patient density (Figure 3: a and b), and finitely similar to work carried out by Singh *et al.*, (2001), Pendse and Singh (1986), Kumar *et al.* (2003) in other geological setting.

The recent study in Yavatmal district dominated by basaltic lithology revealed that groundwater of the area is of bicarbonate (HCO₃⁻) type and high fluoride (F⁻) concentration is observed in deeper aquifers compared to shallow aquifers (Madhnure *et al.*, 2007). Pawar *et al.*, (2008) reported the fluoride ingress in some dug wells exceeding the permissible limit (1 mg/L) of WHO from semi-arid DVP.

5.4 Fluoride Accumulation in Basaltic Groundwater in Drought Situation

Fluoride concentration increases in basaltic groundwater as the result of salinization during drought situation. This is due to the dissolution of secondary minerals in vadose zone. The fluoride concentration increasing above the 1 mg/L (WHO drinking water permissible limit) as the result of low amount rainfall (< 500 mm) is the alarming sign of provoking fluoride related illnesses. However, consecutive drought years (Section 4.2) shows the cumulative built up of fluoride in groundwater from study area as the result of less water available for flushing of accumulated salts in vadose zone.

It is interesting to note that fluoride is present in the urolith composition in the form of CaF₂ (Kale *et al.*, 2011). In north India the relationship between fluoride concentration in drinking water and urolith prevalence in local population is reported by Singh *et al.* (2001). The fluoride is most reactive anionic species it actively takes parts in many human physiological processes. At higher level concentration, sometimes it destroys the cell from various tissues from internal organs. This gives substantial pressure on renal system to filter out protein material of dead cell matter and possibly become nucleus in process of urolith formation (Rotily *et al.*, 2000).

5.5 Role of Fluoride in the Formation of Uroliths

Some dental fluorosis cases are also observed in study area at the time of drought situations. However, no reports are available for skeleton fluorosis in study area. After integrating the entire results, the present study highlights the relationship between excess fluoride ingestion and urolith formation in the population from study area. Fluoride concentration in groundwater also directly affects nervous system in human beings (Alarcon-Herrera *et al.*, 2001) and is linked with cell mortality rate in blood. This can allow higher oxalate generation due to process of dead cell protein breakdown, thus leading to nucleus for urolith (Alarcon-Herrera *et al.*, 2001). However, more research is required from medical researchers in biochemistry point of view to understand the role of fluoride consumption to develop mass urolith epidemics. However, Susheela (1999) reported the other fluoride sources from diet ingredients may be one of the fluoride source.

Fluoride is essential for human in trace concentration to prevent dental caries, while exceeded concentration affects the teeth, bone and nervous system. The deposition of fluoride in the form of calcium fluoride was observed in the uroliths from Saswad (Kale *et al.*, 2011). This indicates the role of fluoride in the formation as well as deposition in concentric layers of uroliths. Anasuya (1982) has demonstrated the relation between high fluoride intake and development of uroliths in rats. Vahlensieck (1985) has been noted the fluoride endemic area experiences the urolithiasis in Europe. However, Singh *et al.* (2001) suggested the proposed mechanism of fluoride participation of in the formation of uroliths. In his view, fluoride is indirectly increasing oxaluria by enhancing the absorption of oxalate from the intestine due to low availability of calcium.

There is no alternative of groundwater for drinking and cooking purpose for people residing in study area. In drought period, they have to be dependent on low yielding local aquifers. As fluoride concentration is increased due to cumulative accumulation triggered by insufficient rainfall, people are use this poor quality water directly for drinking purpose without physical or chemical treatment. As the result of this, consumption of fluoride rich groundwater has enhanced the urolithiasis in local population from year 2000 (Figure 3). This was evidenced by sudden increase of urolith patients as the drought strikes the study area (Figure 3). This necessitates the viable groundwater treatments before used as a drinking as well as cooking purposes to avoid such urolithiasis outbreaks.

5.6 Remedial Measures for Groundwater Fluoride Induced Urolithiasis

To avoid extra ionic load on metabolism water must be used with optimum mineral concentration (Sierakowski *et al.*, 1979). The deionisers present in the market are found to be useful for this purpose. The use of surface sources for drinking e.g. river water or reservoir water

need to undergo regular water disinfection with chlorination, Ultra-violet (UV) light treatment and reverse osmosis (RO). Especially, groundwater has to be filtered through reliable deionising unit before using it for drinking and cooking purposes. Groundwater from study area was contaminated due to considerable amount of fluoride that needs advanced treatments of defluoridation unit. The low cost treatments like using of alum and natural zeolites can be used for removing fluoride ions from groundwater. In low rainfall areas, rainwater is trapped with the help of roof top rainwater harvesting and the collected water can be used for drinking purposes with reliable treatments. With water treatment, study also warrants awareness of local people for drought condition and urolithiasis threats and further management of groundwater fluoride induced urolithiasis from semi-arid parts of DVP area.

6 CONCLUSIONS

The conclusion was drawn from present study that fluoride accumulation is observed during drought period and the collective drought effect was observed on health of local inhabitants in the form of urolithiasis epidemics in the study area. Fluoride concentration exceeded 1 mg/L (WHO permissible limits) in 7 years from 1994 to 2008. The fluoride accumulation in groundwater is observed during drought condition and at the same time urolithiasis cases was reported in substantial number. During drought conditions, the increase in number of hot days in a year were reported and convincingly correlated with urolithiasis occurrence. This highlights the role of climatic conditions on the fluoride ingress in groundwater from study area. Fifteen year groundwater fluoride data is strongly correlated to urolithiasis occurrence. Further, village wise study of fluoride concentration in groundwater and urolithiasis epidemics shows positive correlation. In post-monsoon period has shown fluoride accumulation in groundwater due to dissolution of primary as well as secondary minerals in subsurface aqueous system. In conclusion, present study highlighting the role of excess fluoride consumption through local groundwater system is responsible for increase in urolithiasis from semi-arid Karha basin. The present study warrants the proper groundwater de-fluoridation during drought period to combat urolithiasis occurrence in semi-areas of Deccan Volcanic Province.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

Authors are thankful to University Grants Commission, New Delhi for financial assistance in the form of JRF. Thanks are due to Head, Department of Environmental Sciences, Savitribai Phule Pune University, Pune for providing necessary facilities. Help rendered by Mr. Satyajit Gaikwad during field is thankfully acknowledged. Authors are thankful to anonymous reviewer's suggestion for improving the quality of manuscript.

ABBREVIATIONS

DVP: Deccan Volcanic Province; **GSDA:** Groundwater Surveys and Development Authority; **IMD:** India Meteorological Department; **RO:** Reverse Osmosis; **UV:** Ultra-violet; **WHO:** World Health Organization.

REFERENCES

- A.P.H.A., [American Public Health Association], A.W.W.A. [American Water Works Association] and W.P.C.F. [Water Pollution Control Federation], 1995. *Standard Methods for the Examination Water and Wastewater*. 17th Edition Washington DC, USA.
- Alarcon-Herrera, M. T., Martin-Dominguez, I. R., Trejo-Vazquez R. and Rodriguez-Dozal, S., 2001. Well water fluoride, dental fluorosis, and bone fractures in the Guadiana valley of Mexico. (2001 Research Report 139), *Fluoride*, 34 (920), 139-149.
- Anasuya, A., 1982. Role of Fluoride in Formation of Urinary Calculi. *Studies in Rats. J. Nutr.*, 112, 1787-1795.
- Asper, R., 1984. Epidemiology and socioeconomic aspects of urolithiasis. *Urol Res*, 12, 1-5.
- Costa-Bauza, A, Ramis, M, Montesinos, V, Conte, A, Piza, P, Pieras, E and Grases, F., 2007. Type of renal calculi: variation with age and sex. *World Jour of Urol*, 25, 415-421.
- Chandrajith, R., Wijewardana, G., Dissanayake, C. B. and Abeygunasekara, A., 2006. Biominerology of human urinary calculi (kidney stones) from some geographic regions of Sri Lanka. *Environ. Geochem. and Health*, 28, 393-399.
- Gadgil, A. and Dhorade, A., 2005. Temperature trends in twentieth century at Pune, India, *Atmos Environ*, 39, 6550-6556.
- Handa, B. K., 1975. Geochemistry and genesis of fluoride containing groundwaters of India. *Ground Water*, 13 (3), 275-281.
- Hassan, I., Juncos, L. A., Milliner, D. S., Sarmiento, J. M. and Sarr, M. G., 2001. Chronic renal failure secondary to oxalate nephropathy: a preventable complication after Jejunoileal bypass. *Mayo. Clin. Proc.*, 76, 758-760.
- Indridason, O. S., Birgisson, S., Edwardsson, V. O., Sigvaldason, H., Sigfusson, N. and Palsson, R., 2006. Epidemiology of kidney stones in Iceland: A population-based study. *Scand J Urol Nephrol*, 40, 215-220.
- Kale, S. S., Kadam, A. K., Suyash Kumar and Pawar, N. J., 2010. Evaluating pollution potential of leachate from landfill site, from Pune Metropolitan city and its impact on shallow basaltic aquifers. *Environ Monit and Assess*, 162, 327-346.
- Kale, S. S., Pawar, N. J., Wagholikar, D. S. and Hema A., 2011. Microstructure and growth band studies of uroliths using optical and scanning electron microscopy. *Curr. Sci.*, 100(2), 225-229.
- Kale, S. S. Ghole, V. S., Pawar, N. J. and Jagtap, D. V., 2014. Inter-annual variability of urolithiasis epidemic from semi-arid part of Deccan Volcanic Province, India: climatic and hydrogeochemical perspectives, *International Journal of Environmental Health Research*, 24(3), 278-289.
- Kumar, R., Kapoor, R., Mittal, B., Kumar, A. and Mittal, R. D., 2003. Evaluation of urinary abnormalities in urolithiasis patients: a study from north India. *Ind Jour of Clin Biochem*, 18 (2), 209-215.

- Loughnan, M. E., Nicholls, N. and Tapper, N. J., 2010. The effects of summer temperature, age and socioeconomic circumstance on Acute Myocardial Infarction admissions in Melbourne, Australia. *Int. J. of Health Geographics*, 9, 41.
- Madhavan, N. and Subramanian, V., 2001. Fluoride concentration in river waters of south Asia. *Curr. Sci.*, 80 (10), 1312-1319.
- Madhnure, P., Sirsakar, D. Y., Tiwari, A. N., Ranjan, B. and Malpe, D. B., 2007. Occurrence of fluoride in the groundwaters of Pandharkawada area, Yavatmal District, Maharashtra, India. *Curr. Sci.*, 92, 675-679.
- Muralidharan, D., Nair, A.P. and Satyanarayan, U., 2002. Fluoride in shallow aquifers in Rajgarh Tehsil of Churu District, Rajasthan – an arid environment. *Curr. Sci.*, 83 (6) 25, 699-702.
- Ozsvath, D. L., 2009. Fluoride and environmental health: a review. *Rev. Environ Sci. Biotechnol*, 8, 59-79.
- Pawar, N. J., Pawar, J. B., Kumar S. and Supekar A., 2008. Geochemical eccentricity of ground water allied to weathering of basalt from the Deccan volcanic province, India: Insinuation on CO₂ consumption. *Aqua Geochem*, 14, 41-71.
- Pendse, A. K. and Singh, P. P., 1986. The Etiology of Urolithiasis in Udaipur (Western Part of India). *Urol Res*, 14, 59-62.
- Rotily, M., Leonetti, F., Iovonna, C., Berthezene, P., Dupuy, P., Vazi. A. and Berland, Y., 2000. Effects of low animal protein or high-fiber diets on urine composition in calcium nephrolithiasis. *Kid. Int.*, 57, 1115-1123.
- Sierakowski, R., Finlayson, B. and Landes, R., 1979. Stone Incidences as related to water hardness in different geological regions of the United States. *Urol. Res.*, 7, 157-160.
- Singh, P. P., Barjatiya, M. K. and Dhing, S., 2001. Evidence suggesting that high intake of fluoride provokes the nephrolithiasis in tribal population. *Urol. Res.*, 29, 238-244.
- Susheela, A. K., 1999. Fluorosis management programme in India. *Curr. Sci.* 77 (10), 1250-1256.
- Vahlensieck, W., 1985. Influence of water quality on urolithiasis. In: Schuille PO, Smith LH, Robertson WG, Vahlensieck W (eds) *Urolithiasis and related research*. Plenum, New York, 97.
- Whitford, G. M., 1996. The metabolism and toxicity of fluoride. *Monographs in oral science*, 2nd rev. edⁿ, volume 16. Karger, New York.
